

Heterogeneous and metasomatized mantle recorded by mineral trace elements in Donghai garnet peridotites of the Sulu UHP terrane, eastern China

J. ZHENG^{1,2,3}, R.Y. ZHANG², J.G. LIOU³, W.L. GRIFFIN^{3,4} AND S.Y. O'REILLY³

¹ Earth Sciences, Univ. of Geosci., Wuhan 430074, China

² Geol. and Env. Sci., Stanford University, CA 94305, USA

³ GEMOC ARC National Key Centre, Macquarie University, NSW 2109, Sydney, Australia

⁴ CSIRO Explor. and Mining, NSW 1067, Sydney, Australia

Garnet peridotites from the Sulu terrane of eastern China were derived from the mantle wedge above a subduction zone of the Yangtze craton beneath the Sino-Korean craton. They underwent Triassic ultrahigh-pressure (UHP) metamorphism with their country rock gneisses. Mineral trace elements of Donghai garnet peridotites were analyzed by laser ablation ICPMS. One Zhimafang peridotite (porphyroblastic texture) consists of olivine ($Fo_{91.2-91.8}$), enstatite ($En_{92.0}$), garnet ($Prp_{66.5-70.7}$) and diopside ($Di_{90.8}$). Four Xugou samples contain nearly equigranular olivine ($Fo_{90.6-92.2}$), enstatite ($En_{92.0-92.7}$), garnet ($Prp_{63.2-87.1}$) and diopside ($Di_{93.7-95.7}$) with additional minor phlogopite. Garnets from both localities have low LREE and show negative Ce anomalies, but the Zhimafang garnet has higher HREE (8-15) than those from Xugou (2-6). Zhimafang diopside shows sinusoidal REE pattern with flat HREE. Xugou diopsides, in contrast, have LREE-enriched patterns with large variation in HREE. Diopsides from both peridotites have low abundance of Yb, Y and Ti, and low Ti/Eu (517-1158), but high and variable Sr/Nd (4800-22000) and $(La/Yb)_n$ (40-100). Geochemical modeling indicates that the concentrations of moderately incompatible trace elements such as Y, Ti, Yb and Dy in diopside fit a fractional melting trend with about 30% partial melting of a primitive mantle source. However, Zr and Gd in the Zhimafang diopside, and the highly incompatible trace elements (such as Nb and especially LREE, Sr, Th and U) of diopside from both bodies, cannot be described by this model, and are attributed to carbonatitic metasomatism. The Sr isotopic composition of diopside and garnet, and marked negative Ce anomalies of garnets, suggest a sedimentary fluid source. Hydrous silicates (eg. Phlogopite) and low whole rock CaO/Al_2O_3 suggest a later silicate metasomatism related to subducted crustal materials. Systemic differences in the Ni content and Mg of olivine, and the Y, Al, HREE, Nd/Y and Sc/Y of garnet, show that the mantle volume represented by the Zhimafang peridotite is less depleted than the Xugou mantle. Such differences indicate that the garnet peridotites derived from a heterogeneous mantle in which different volumes had different evolution histories prior to their involvement in Triassic continental subduction.