Enriched mantle component in high Sr/Y ("TTG") granites: Hf, Sr, Nd, O, isotopic compositions of cretaceous arc magmas from New Zealand

A.J. Tulloch[1], W.L. Griffin[2], D.L. Kimbrough[3], K. Faure[4], A. Saeed[2]

[1]GNS Science, Dunedin, New Zealand (a.tulloch@gns.cri.nz)
[2]Macquarie University, Sydney, Australia (wgriffin@els.mq.edu.au)
[3]San Diego State University, CA, USA (dkimbrou@geology.sdsu.edu)
[4]GNS Science, Lower Hutt, New Zealand (k.faure@gns.cri.nz)

High Sr/Y plutonic rocks (HiSY, TTG, adakitic) form significant components of continental margin magmatism. Some are associated with major flare ups in rates of magmatism. Jurassic-Early Cretaceous subduction along the NZ segment of east Gondwana produced margin-parallel paired belts over at least 800 km: an inboard high Sr/Y (>40) SPS belt (130–105 Ma) immediately postdates an outboard, low Sr/Y, Darran belt (170–128 Ma) (Tulloch and Kimbrough, 2003).

A Sr–Nd isotope array for SPS (McCulloch et al., 1987) extending steeply below the mantle array from the near-DM Darran field (eNd = +3.5 to 0.7) was interpreted as including a low-Rb/Sr basaltic component with a significant but limited prehistory (?Darran belt). Partial melting of an underplated and/or underthrust Darran belt has also been suggested to explain the high Sr/Y, and the inboard location (McCulloch et al., 1987; Muir et al., 1995; Tulloch and Kimbrough, 2003).

We report new Hf isotope data for SPS (eHf = 3.9–7.5) that require a more enriched component, with lower eHf, in addition to a Darran component (eHf = 9.8) (or oceanic crust/ lithosphere). Qtz, zircon & opx ^{18}O (calculated WR = 4.8–7.6‰) and feldspar Pb isotopic data rule out significant continental crust, and pelagic sediment (low Sr_i rules out EMII), pointing instead to EMI enriched mantle. A significant mantle component is also required to supply heat to drive the observed SPS flare up.

References

McCulloch, M.T. et al., 1987. *CMP* **97**, 183–195.
Muir, R.J. et al., 1995. *Jl Geol. Soc.* **152**, 689–701.
Tulloch, A.J., Kimbrough, D.L., 2003. *Geol. Soc. Am. Special Paper* **374**, 275–295.