

Vegetation over hydrologic control of sediment transport over the past 100,000 yr

A. DOSSETO¹, S.P. TURNER¹, P. HESSE², K. MAHER³ AND K. FRYIRS²

¹GEMOC National Key Centre, Department of Earth and Planetary Sciences, Macquarie University, Sydney, Australia (adosseto@els.mq.edu.au)

²Department of Physical Geography, Macquarie University, Sydney, Australia

³Department of Geological and Environmental Sciences, Stanford University, USA

Uranium isotopes can be used to determine the residence time of sediments in a catchment, i.e. how long they are stored in weathering profiles and transported through the catchment by rivers. We have measured uranium isotopes in sediments from palaeo-channels of the Murrumbidgee River (Murray-Darling Basin, southeastern Australia) to quantify variations in sediment residence times over the past 100,000 years.

Results indicate that sediments transported through the Murrumbidgee catchment during the Last Glacial Maximum (LGM) resided for 10's of thousands of years in the catchment. This contrasts with modern and 100ka-old channel sediments where the residence time reaches values as high as 400,000-500,000 years. Variations in sediment residence time in the Murrumbidgee basin do not strictly follow changes in bankfull discharge but instead are correlated with shifts in vegetation and atmospheric CO₂. In the absence of significant glacial erosion in this basin during LGM, this is at odds with what is expected from the links between climate and erosion (a decrease in CO₂ and temperature is expected to induce a decrease in weathering and erosion). Vegetation may be the link between climate and sediment transport: sparse vegetation in the upper catchment allows significant hillslope erosion during LGM but dense woodlands in the Holocene and during the last interglacial inhibit sediment delivery to the river from hillslopes and sediments are derived from the re-working of old (a few 100s ka) alluvial deposits. These observations would suggest that (i) changes in hydrology cannot explain alone changes in sediment transport and (ii) the impact of climate change on catchment erosion is operating indirectly, via changes in vegetation type and density.